

Patient Medication Information

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr **DIPRIVAN® 1%**

Propofol

This patient medication information is written for the person who will be taking **DIPRIVAN 1%**. This may be you or a person you are caring for. Read this information carefully. Keep it as you may need to read it again.

This patient medication information is a summary. It will not tell you everything about this medication. If you have more questions about this medication or want more information about **DIPRIVAN 1%**, talk to a healthcare professional.

What DIPRIVAN 1% is used for:

DIPRIVAN 1% is used:

- in adults and children 3 years of age and older during surgery and other medical procedures to make the patient become unconscious (asleep).
- in adults in the Intensive Care Unit (ICU) and in other medical settings to keep patients sedated (a very relaxed state of calm, drowsiness or sleep).

How DIPRIVAN 1% works:

DIPRIVAN 1% belong to a group of medicines called general anesthetics. These types of medicines work by causing a loss of feeling and awareness that feels like being in a deep sleep.

The ingredients in DIPRIVAN 1% are:

Medicinal ingredient(s): propofol

Non-medicinal ingredients: disodium edetate, egg phosphatide, glycerol, sodium hydroxide, soybean oil and water for injection.

DIPRIVAN 1% comes in the following dosage form(s):

Emulsion; 10 mg / mL

Do not use DIPRIVAN 1% if:

- In children under the age of 18 years for sedation during surgical/diagnostic procedures or in the intensive care unit (ICU).
- If you have ever received DIPRIVAN 1% before and have experienced an allergic reaction to its use or if you know that you are allergic to propofol or any of the nonmedicinal ingredients (including eggs or egg products and soybeans or soy products) in DIPRIVAN 1% (see **What are the ingredients in DIPRIVAN 1%**)

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take DIPRIVAN 1%. Talk about any health conditions or problems you may have, including if you:

- All health problems you have now or have had in the past.
- Other medicines you take, including ones you can buy without a prescription.
- If you have any other health problems such as problems with your:
 - heart,
 - pancreas,
 - thyroid,
 - breathing,
 - kidneys or liveror if you have been generally unwell for some time.
- If you have ever had an epileptic fit or seizures.
- If you have ever been told that either you have very high fat levels in your blood or your body has problems in breaking down fat adequately.
- If you have diabetes.
- If you are pregnant, plan to become pregnant or are breastfeeding.

Other warnings you should know about:

Driving and using machines: DIPRIVAN 1% may temporarily interfere with your ability to drive or operate tools or machinery. You should not drive or operate machinery until all the effects of DIPRIVAN 1% have worn off.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with DIPRIVAN 1%:

- Alfentanil (used as painkillers for initiating and/or maintaining general anaesthesia).
- Dexmedetomidine (used to induce sedation (a very relaxed state of calm, drowsiness or sleep))
- Sevoflurane (used to induce and maintain general anaesthesia)
- Midazolam (used to induce sedation (a very relaxed state of calm, drowsiness or sleep) and relieves anxiety and muscle tension).
- Opioid medicines, such as fentanyl (used to treat severe pain)
- Other medicines that have an effect on the nervous system, breathing and blood pressure.
- Avoid alcohol before and for at least 8 hours after receiving DIPRIVAN 1%.

How to take DIPRIVAN 1%:

DIPRIVAN 1% will be given to you by your anaesthesiologist or intensive care doctor.

Usual dose:

The dose given is decided by the healthcare professional based on the medical procedure it is being used for and your physical condition.

When Receiving DIPRIVAN 1%

DIPRIVAN 1% will be given to you as an injection into a vein, usually in the back of the hand or in the forearm. Your anaesthesiologist may use a needle. For long operations and for use in intensive care situations, an electric pump may be used to control the rate at which the injection is given.

Your anaesthesiologist or intensive care doctor will closely control the amount of DIPRIVAN 1% that is given to you. The amount will be adjusted according to how deeply your anaesthesiologist or intensive care doctor wishes you to sleep or be sedated. He/she will also take account of your age and physical fitness and adjust the amount accordingly.

Several different medicines may be needed in order to keep you asleep or sedated, free from pain, breathing in a healthy way and to keep your blood pressure steady. Your anaesthesiologist or intensive care doctor will decide which medicines to use as the need arises.

If DIPRIVAN 1% is used for a long time in intensive care situations, certain patients may need to be given zinc (a mineral) supplements.

After Receiving DIPRIVAN 1%

You may feel some pain in the arm into which DIPRIVAN 1% is given; this is normal.

Overdose:

If you think you, or a person you are caring for, have taken too much DIPRIVAN 1%, contact a healthcare professional, hospital emergency department, regional poison control centre or Health Canada's toll-free number, 1-844 POISON-X (1-844-764-7669) immediately, even if there are no signs or symptoms.

Possible side effects from using DIPRIVAN 1%:

These are not all the possible side effects you may have when taking DIPRIVAN 1%. If you experience any side effects not listed here, tell your healthcare professional.

Serious side effects and what to do about them

Frequency/Side Effect/Symptom	Talk to your healthcare professional		Stop taking the/this drug (if applicable) and get immediate medical help
	Only if severe	In all cases	
During use of DIPRIVAN 1%			
Very common			
A feeling of pain near the injection site	✓		
Common			
A fall in blood pressure		✓	
A slower heart beat		✓	
Changes in your breathing pattern		✓	
Rare			
Some twitching and shaking		✓	
Feeling of euphoria (sense of happiness)		✓	
Very rare			
Allergic reactions: swelling of face, hives, rash and itchiness, including anaphylactic shock and anaphylactoid reaction			✓
Fluid in the lungs: severe shortness of breath, difficulty breathing and wheezing, especially when lying down			✓
Unknown			
Discolouration of the urine		✓	
Breakdown of muscle cells (rhabdomyolysis): muscle pain, muscle weakness and dark urine			✓
After use of DIPRIVAN 1%			
Common			
Nausea and vomiting	✓		
Headache	✓		
Uncommon			
Redness or soreness where DIPRIVAN 1% was given	✓		
Rare			
Some twitching and shaking		✓	

Frequency/Side Effect/Symptom	Talk to your healthcare professional		Stop taking the/this drug (if applicable) and get immediate medical help
	Only if severe	In all cases	
Feeling of euphoria (sense of happiness)		✓	
Very rare			
Inflammation of the pancreas (pancreatitis): long periods of pain in the stomach and/or intestine area may go around to your back. You may also vomit		✓	
Fluid in the lungs: severe shortness of breath, difficulty breathing and wheezing, especially when lying down		✓	
Feeling hot, have increased body temperature or fever, or flushed skin		✓	
A feeling of sexual arousal or long-lasting penile erection		✓	
Fainting or loss of consciousness		✓	
Unknown			
Hepatitis (inflammation of the liver), acute liver failure (symptoms can include yellowing skin and eyes, itching, dark coloured urine, stomach pain and liver tenderness (indicated by pain under the front of the rib cage on your right-hand side), sometimes with loss of appetite)		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting side effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your healthcare professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Keep out of reach and sight of children.

If you want more information about DIPRIVAN 1%:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes the Patient Medication Information by visiting the Health Canada Drug Product Database website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website www.aspenpharma.ca; or by calling 1-844-330-1213.

This leaflet was prepared by

Aspen Pharmacare Canada Inc.
201 - 2030 Bristol Circle
Oakville, ON, L6H 0H2

Date of Authorization: 2025-07-28

Trademarks are owned by or licensed to the Aspen Group of companies.
©2025 Aspen Group of companies or its licensor. All rights reserved.