

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE
PATIENT MEDICATION INFORMATION

EMLA® Cream
Lidocaine 2.5% and Prilocaine 2.5% Cream

Read this carefully before you start taking **EMLA® Cream** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **EMLA® Cream**.

What is EMLA® Cream used for?

EMLA Cream is used to temporarily numb the skin. It can be used:

- on healthy, unbroken skin before getting a needle or having blood taken;
- on healthy, unbroken skin before minor skin procedures or surgeries;
- before being vaccinated with only the following vaccines:
 - MMR (Measles-Mumps-Rubella)
 - DPTP (Diphtheria-Pertussis-Tetanus-Poliovirus)
 - *Haemophilus influenzae b*
 - Hepatitis B.
- on the genitals for short surgeries;
- for the cleansing of leg ulcers.

EMLA Cream should only be used for the uses above. Serious and life-threatening side effects have happened when EMLA Cream was applied to large areas of skin.

How does EMLA® Cream work?

EMLA Cream contains the drugs lidocaine and prilocaine. These drugs are anesthetics and they cause a temporary numbness of the skin where the cream is applied.

What are the ingredients in EMLA® Cream?

Medicinal ingredients: lidocaine 2.5% and prilocaine 2.5%

Non-medicinal ingredients: carboxypolymethylene, polyoxyethylene hydrogenated castor oil, sodium hydroxide, and purified water.

The 5 g EMLA Cream tube also includes dressings. Tegaderm® dressings contain polyether polyurethane films, acrylate adhesives and paper liners. These dressings are hypoallergenic and do not contain latex.

EMLA® Cream comes in the following dosage forms:

Cream: 5 g and 30 g tubes

Do not use EMLA® Cream if:

- you are allergic to lidocaine, prilocaine, any other “-caine” type anesthetics, or any of the non-medicinal ingredients in the product.
- you have a blood disorder called methemoglobinemia;
- it is for infants who are 12 months of age or younger and are taking drugs that may cause the blood disorder called methemoglobinemia (e.g., sulphonamides);
- it is for infants who were born after less than 37 weeks of pregnancy.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take EMLA[®] Cream. Talk about any health conditions or problems you may have, including if you:

- have glucose-6-phosphate dehydrogenase deficiency;
- have ever had a bad, unusual or allergic reaction to lidocaine or prilocaine, also available under brand names such as Xylocaine[®] (lidocaine) and Citanest[®] (prilocaine);
- may be sensitive or allergic to other ingredients of the cream or Tegaderm[®] dressing;
- have an infection, skin rash or cut at, or near, the area where you want to apply EMLA Cream;
- have atopic dermatitis, eczema or any other skin problems or diseases;
- have severe kidney or liver disease;
- are pregnant, trying to become pregnant or are breastfeeding;
- would like to use EMLA Cream prior to treatment of a leg ulcer(s);
- would like to use EMLA Cream on the genital area of children.

Other warnings you should know about:

- **Risk of serious side effects**

Be careful to apply no more than the maximum recommended dose of EMLA Cream. Serious and life threatening side effects have occurred when EMLA Cream was not used properly. This includes a serious blood disorder called methemoglobinemia, where the blood brings less oxygen to your body. If you develop this condition, you may have brownish or greyish skin especially around lips and nails. If your condition worsens, you can have:

- Heart problems leading to a heart attack, difficulty breathing or not breathing
- Problems with your central nervous system such as:
 - Seizures
 - Dizziness
 - Fainting,
 - Feeling drowsy

Other symptoms you may have include:

- numbness of the tongue,
- light-headedness,
- confusion,
- headache,
- sight or hearing problems,
- vomiting,
- dizziness,
- weakness,
- nervousness,
- unusual sweating,
- trembling

If you get any of these side effects, stop taking EMLA Cream and get medical help right away.

- Children are at greater risk for serious side effects. Always follow your doctor's instructions for using EMLA Cream, especially in young children and infants under 3 months old. EMLA Cream should not be used on the genitals of children or infants.

- EMLA Cream should not be used close to the ear or eyes because it can cause damage.
- Know how you feel after using EMLA Cream before you drive or use heavy machines.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with EMLA® Cream:

- drugs for heart rhythm problems (e.g. mexilitine, amiodarone);
- other anesthetics;
- other drugs which may cause methemoglobinemia, including: sulfonamides, acetanilide, aniline dyes, benzocaine (or other “-caine” type anesthetics), chloroquine, dapsone, naphthalene, nitrates or nitrites, nitrofurantoin, nitroglycerin, nitroprusside, pamaquine, para-aminosalicylic acid, phenacetin, phenobarbital, phenytoin, primaquine, quinine and high doses of acetaminophen.


How to take EMLA® Cream:

- Be careful not to apply more EMLA Cream than your doctor recommended. Serious and life threatening side effects have occurred when EMLA Cream was not used properly and more than the recommended amounts were used.
- Be sure to only put EMLA Cream on unbroken skin, except for leg ulcers where it has been prescribed for that condition.
- Do not put EMLA Cream in your mouth, or swallow it. If EMLA Cream is accidentally swallowed, call your doctor.
- Do not re-use EMLA Cream dressings once applied.
- The numbing effect of EMLA Cream starts working about 1 hour after it is applied. You may still feel pressure and touch in the area where you applied EMLA Cream. The numbness of the skin may continue to increase after the cream is removed and will last for at least 2 hours following a 1-2 hour application.
- Tell your doctor or pharmacist if you feel that the effect of EMLA Cream is too strong or too weak.

Steps to apply EMLA Cream:




1. Make sure your skin is clean and dry. Apply cream in a thick layer at the site of the procedure. Do **not** rub the cream into the skin.




2. Cover treated area with an air-tight dressing such as Tegaderm® or plastic wrap. Tegaderm® is provided with the 5 g tubes only. If using Tegaderm® remove the center cut-out piece as shown. Peel the paper liner from the paper-framed dressing. It is important to cover EMLA Cream with an air-tight dressing to ensure that the cream penetrates the skin properly and numbness of the area is felt.



3. Carefully cover the EMLA Cream so that you are left with a thick layer of cream underneath the dressing. Do not spread out the cream. Smooth down the dressing edges carefully and make sure it is secure to prevent leakage. If using plastic wrap, hold the dressing in place with adhesive or medical tape and make sure it is air-tight.



4. If using Tegaderm®, remove the paper frame. The time of application can easily be marked directly on the Tegaderm® with a ballpoint pen. If using plastic wrap, mark the time of application on the medical tape that is holding the dressing in place.



5. Keep the dressing and cream on for as long as your doctor recommended, usually at least 1 hour. Remove the dressing and clear the area of excess cream thoroughly before the procedure. If you are applying the EMLA Cream for a procedure to be performed by a doctor, you should leave the dressing on for the doctor to remove, unless otherwise instructed.

6. Throw the plastic wrap or the Tegaderm® in the garbage, out of the reach and sight of children and pets.

Usual dose:

- The dosing for adults and children is listed below. To help prevent side effects, your doctor may need to adjust the usual dose if you:
 - are elderly
 - are acutely ill
 - have severe liver or kidney disease
 - are being treated with other anesthetics or certain heart drugs (e.g. mexilitine, amiodarone)
 - have skin conditions such as atopic dermatitis
- When applying EMLA Cream, 1 g of cream is the same as a ribbon of cream of about 3.5 cm (1.5 inches).
- Make sure to follow the steps above when you apply EMLA Cream, including covering the cream with an air-tight dressing.

Adults

DOSAGE OF EMLA CREAM ON HEALTHY SKIN

For minor skin procedures/surgeries or when getting a needle or having blood taken:

- apply a thick layer of cream, about half of a 5 g tube (2 g), on an area slightly larger than a two dollar coin or “toonie”.
- leave the EMLA Cream on for at least 1 hour.

For procedures with larger areas, such as skin grafting:

- Your doctor may apply the EMLA Cream when you arrive for your procedure.
- If your doctor told you to put the cream on yourself before the procedure, make sure they clearly explain the size of the area to be treated:
 - apply a thick layer of cream to the area to be treated, about 1.5 to 2 g of cream to an area that is 10 cm²;
 - 1.5 to 2 g is about half of a 5 g tube;
 - a 10 cm² area is a little larger than the size of a two dollar coin or “toonie”).
 - leave the EMLA Cream on for at least 2 hours.

You will not get any added benefit from leaving EMLA Cream on for longer than 5 hours.

DOSAGE OF EMLA CREAM ON LEG ULCERS

Talk to your doctor **before** using EMLA Cream on leg ulcers.

For topical anesthesia before your leg ulcer(s) are cleansed:

- apply a thick layer of cream to the area to be treated, about 1.5 to 2 g of cream to an area that is 10 cm²;
 - 1.5 to 2 g is about half of a 5 g tube;
 - a 10 cm² area is a little larger than the size of a two dollar coin or “toonie”).
- use no more than 10 g (two 5 g tubes).
- leave EMLA Cream on the leg ulcer(s) for at least 30 minutes. Leaving EMLA Cream on for 60 minutes may improve the numbness you feel.
- the cleansing of the leg ulcer(s) should begin within 10 minutes after removing the cream.
- discard the tube of EMLA Cream with any remaining contents after each treatment.

DOSAGE OF EMLA CREAM ON THE GENITALS

Talk to your doctor **before** using EMLA Cream on your genitals.

Apply EMLA Cream when you are with your doctor because it only needs to be on your skin for 5-10 minutes before the procedure.

For needle insertion or for surgical treatment of small lesions (removal of warts or a biopsy):

- use half of a 5 g tube (2 g) on the area to be treated. You may need to put the cream on a few areas.

You do not need an airtight dressing when using EMLA Cream on the genitals. Your doctor should begin the surgical procedure immediately after removing the cream.

Children

- Observe your child closely during and after use of topical anesthetics. Children are at greater risk than adults for serious side effects, especially infants under 3 months of age.
- When using EMLA Cream for your child's pain relief, remember it is also very important to provide comfort and emotional support.
- In children, EMLA Cream should only be applied to healthy, unbroken skin.

Infants Under the Age of 3 Months:

- Use only for infants under the age of 3 months if instructed by a doctor.
- Apply up to 1 g of cream on a skin area not larger than 10 cm² (a little larger than the size of a two dollar coin or "toonie").
- Leave on for 1 hour. Do NOT leave EMLA Cream on the skin for longer than 1 hour.

Infants Between 3 and 12 Months of Age:

- Apply up to 2 g of cream on a total skin area not larger than 20 cm² (a little larger than the size of a credit card).
- Leave on for at least 1 hour. Do not leave on the skin for more than 4 hours.

Children Between 1-6 Years:

- Apply up to 10 g of cream on a total skin area not larger than 100 cm² (a little larger than the size of two credit cards).
- Leave on for at least 1 hour. Do not leave on the skin for more than 5 hours.

Children Between 7-12 Years:

- Apply up to 20 g of cream on a total skin area not larger than 200 cm² (a little larger than a standard postcard).
- Leave on for at least 1 hour. Do not leave on the skin for more than 5 hours.

Overdose:

<p>If you think you have taken too much EMLA Cream, contact your doctor, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms. You may require medical attention.</p>

What are possible side effects from using EMLA[®] Cream?

These are not all the possible side effects you may feel when taking EMLA Cream. If you experience any side effects not listed here, contact your healthcare professional.

Your skin where EMLA Cream was applied may stay numb for up to several hours after you take the cream off. Be careful to avoid hurting your skin until your skin isn't numb anymore. This includes scratching, rubbing or exposing your skin to extreme hot or cold temperatures.

The following side effects can happen to the skin where the cream was applied:

- whitening or redness
- slight swelling or puffiness
- initial burning or itching
- small red dots or purple spots

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE Methemoglobinemia: brownish or greyish skin especially around lips and nails			X
Eye irritation: if EMLA Patch enters the eye, immediately rinse the eye in water.			X
Allergic reaction: difficulty swallowing or breathing, wheezing; drop in blood pressure; feeling sick to your stomach and throwing up; hives or rash; swelling of the face, lips, tongue or throat.			X

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

<p>Reporting Side Effects</p> <p>You can report any suspected side effects associated with the use of health products to Health Canada by:</p> <ul style="list-style-type: none"> • Visiting the Web page on Adverse Reaction Reporting (https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html) for information on how to report online, by mail or by fax; or • Calling toll-free at 1-866-234-2345. <p><i>NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.</i></p>
--

Storage:

- Keep EMLA Cream well out of the reach and sight of children.
- Store EMLA Cream at room temperature (15-30°C). Protect from freezing.
- Do not use EMLA Cream after the expiry date on the tube or after 30 days of opening the tube. Remember to return any unused EMLA Cream to your pharmacist.

If you want more information about EMLA® Cream:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer's website www.aspenpharma.ca, or by calling 1-844-330-1213.

This leaflet was prepared by
8 – 1155 North Service Road West
Oakville, ON
L6M 3E3

Trademarks are owned by or licensed to the Aspen Group of companies.

Tegaderm® - Registered trademark of 3M.

Last Revised: August 9, 2019